



Crown/Bridge & Inlay/Onlay After Care Instructions

Jason Tanguay, DDS * Lindsey Hollern, DDS

MINT.



dental studio

After Care Instructions:

Some amount of pain and tenderness is normal after a crown or other longer restorative appointment. It is common to have temperature sensitivity sense slight changes in bite after your temporary restoration is placed.

- * Acetaminophen or ibuprofen (or whatever you would use for a headache) may be taken every 4-6 hours for tenderness or soreness.
- * Be gentle to this area and avoid hard or chewy foods to provide adequate healing time.
- * If temporary restoration is displaced and there is no sensitivity,, this is not an emergency. If your appointment to seat your permanant restoration is just a few days away, there is no need to replace the temporary. If you are more than a few days away from your permanant restoration it is important to get a new temorary to hold the tooth in its current position - please call the office to schedule a quick appointment. In some cases we may provide you with temporary cement, if you have temporary cement and your restoration has come off please follow the instructions below:
 - * Clean out residual cement from temporary crown/onlay, etc. and practice fitting on to tooth to ensure proper placement.
 - * Mix equal parts Base and Catalyst temporary cement provided at appointment and apply thin layer to temporary restoration.
 - * Dry tooth and place. Apply firm pressure for 1 ½ minutes.
 - * Pull knotted floss through the space between temporary and neighboring teeth to remove excess cement.